

Neuropeptides in the pathophysiology and treatment of cachexia

Stephanie M. Krasnow and Daniel L. Marks

Department of Pediatrics, Oregon Health & Science University, Portland, Oregon, USA

Correspondence to Dr Daniel L. Marks, Department of Pediatrics, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Mail Code L481, Portland, OR 97239, USA
Tel: +1 503 494 6218; fax: +1 503 494 5235; e-mail: marksd@ohsu.edu

Current Opinion in Supportive and Palliative Care 2010, 4:266–271

Purpose of review

Cachexia occurs in various inflammatory diseases and is characterized by weight loss and muscle wasting. Pro-inflammatory cytokines modulate the activity of neuropeptides and hormones that control energy homeostasis and/or illness behaviors. This review summarizes recent (published within the past 18 months) literature regarding neuropeptides and hormones that have been implicated in the pathophysiology of cachexia, and that are likely to have therapeutic potential for preventing or reversing cachexia in various disease states.

Recent findings

Hypothalamic pro-opiomelanocortin (POMC) and agouti-related protein (AgRP) neurons are downstream targets for pro-inflammatory cytokines. Genetic or pharmacological blockade of melanocortin receptor signaling preserves lean body mass and attenuates anorexia in experimental models of cachexia. Orally available melanocortin receptor antagonists have been developed and tested in cachectic animals with favorable results. Ghrelin and ghrelin mimetics increase appetite and preserve lean body mass in cachectic patients with diverse underlying diseases. Additional neuropeptide-expressing neurons in the hypothalamus (e.g., orexin neurons) might play a role in cachexia-associated lethargy.

Summary

Promising outcomes from recent preclinical studies and/or early clinical trials with melanocortin receptor antagonists and ghrelin mimetics raise hopes that safe and effective anti-cachexia drugs will soon become available for widespread clinical use.

Keywords

cachexia, cytokines, ghrelin, inflammation, melanocortin

Curr Opin Support Palliat Care 4:266–271
© 2010 Wolters Kluwer Health | Lippincott Williams & Wilkins
1751-4258

Introduction

Cachexia, or disease-associated wasting, is a metabolic disorder that has a profound negative impact upon morbidity, mortality, and quality of life. Cachexia occurs in many infectious and chronic diseases (e.g., acquired immunodeficiency syndrome, cancer, congestive heart failure, and chronic kidney disease) [1–4]. Cachexia was recently defined as ‘a complex metabolic syndrome associated with underlying illness and characterized by loss of muscle with or without loss of fat mass’ [5], p. 794). In addition to the characteristic weight loss (in adults) or growth failure (in children), cachectic individuals commonly exhibit anorexia, increased resting energy expenditure, and lethargy. Nutritional intervention alone is insufficient to restore lean body mass in cachectic patients [6], and effective pharmacological treatments that prevent or reverse the symptoms of cachexia have so far remained elusive.

Cytokines and peripheral inflammation

The cause of cachexia is multifactorial, involving complex interactions between immune, metabolic, endocrine, and neural mediators. Inflammation is believed to play a causal role in the pathogenesis of cachexia across diverse disease states [7]. Cachexia is associated with elevated circulating levels of pro-inflammatory cytokines [e.g., interleukin-1 β (IL-1 β), interleukin-6, tumor necrosis factor- α (TNF- α), and leukemia inhibitory factor (LIF)] in humans [8,9]. Pro-inflammatory cytokines, which are released by peripheral immune cells in response to infection or tissue damage, act upon their target cells in a paracrine or endocrine manner. Although pro-inflammatory cytokines can promote catabolism via direct effects on skeletal muscle and adipose tissue *in vitro* [10,11], many of the metabolic and behavioral actions of pro-inflammatory cytokines involve cytokine signaling within the central nervous system (CNS).

Cytokines and central inflammation

Systemic inflammatory responses are amplified by the local production of pro-inflammatory cytokines within the CNS [12]. These brain-derived cytokines act upon neural circuits in the hypothalamus and brainstem that modulate energy homeostasis, hormone secretion, and autonomic function [13[•]]. Acute inflammatory stimuli activate nuclear factor kappa B (NF- κ B, a transcription factor upon which many inflammatory signaling pathways converge) [14] and induce expression of the immediate early gene *c-fos* (a marker of neuronal activation) [15] in discrete hypothalamic and brainstem nuclei. The bacterial endotoxin lipopolysaccharide (LPS) elicits profound anorexia in wild-type mice; however, a sustained LPS-induced reduction in feeding is not observed in mice with genetically disrupted inflammatory signaling pathways, not even in animals that are transplanted with wild-type circulating immune cells [16]. Collectively, these observations suggest that central inflammation plays a critical role in illness-induced pathophysiology.

In rodents, intracerebroventricular (ICV) cytokine administration recapitulates many of the features of cachexia [17–20]. Rats that received daily ICV injections of TNF- α for 4 days exhibited reduced food intake and body mass, increased oxygen consumption, and enhanced brown adipose tissue thermogenesis compared to saline-treated animals [21[•]]. Blockade of hypothalamic TNF- α signaling with infliximab increased food intake in tumor-bearing rats, partially restored body weight in rats with experimentally induced sepsis, and improved survival in both experimental models of cachexia, thus lending support to the hypothesis that central cytokine signaling plays an obligatory role in cachexia.

Many neuropeptide-expressing neurons in the hypothalamus and brainstem participate in the control of feeding and metabolism. Some of these same neurons are responsive to inflammatory challenges. This review will highlight some of the neuropeptides and peptide hormones that modulate energy homeostasis and/or sickness behaviors, and that have also been implicated in the pathophysiology of cachexia. More importantly, many of these neural/endocrine mediators have been identified as promising therapeutic targets for the prevention and treatment of cachexia of various etiologies.

Melanocortins

The hypothalamic arcuate nucleus (ARC) is an important target for pro-inflammatory cytokines. The ARC contains two anatomically distinct populations of neurons that have opposing effects on energy homeostasis. One set of neurons produces α -melanocyte-stimulating hormone

(α -MSH), an anorexigenic neuropeptide that is derived from the proopiomelanocortin (POMC) precursor. POMC neurons also reside in the nucleus of the solitary tract in the brainstem. α -MSH inhibits feeding and increases energy expenditure by activating melanocortin receptors, primarily the type 4 melanocortin receptor (MC4-R) [22]. A second population of neurons in the ARC secretes the orexigenic peptides neuropeptide Y (NPY) and agouti-related protein (AgRP). AgRP stimulates feeding and decreases metabolism by acting as an inverse agonist at the MC4-R [23].

POMC and AgRP neurons in the ARC are responsive to pro-inflammatory cytokines. Both POMC and AgRP neurons express the type I interleukin-1 receptor, and IL-1 β stimulates α -MSH secretion and inhibits AgRP release from hypothalamic explants [24,25]. Interleukin-7 increases POMC mRNA in fed mice and reduces AgRP gene expression in mice that are re-fed for 4 h following an overnight fast [26[•]]. Grossberg *et al.* [27^{••}] recently demonstrated that POMC neurons are direct targets for central LIF signaling. ARC POMC neurons express LIF receptor mRNA and are transcriptionally activated by central LIF treatment. Acute ICV LIF injection transiently suppresses food intake in wild-type mice and stimulates α -MSH secretion from wild-type hypothalamic explants. Transgenic mice that lack gp130 (the signal transducing subunit of the LIF receptor complex) only in POMC neurons exhibit attenuated LIF-induced α -MSH secretion and lack an anorectic response to ICV LIF treatment. Thus, intact LIF receptor signaling in POMC neurons is necessary for acute LIF-induced anorexia in mice. It remains to be determined whether a sustained activation of POMC neurons by chronically elevated LIF (or IL-1 β) concentrations in the brain is necessary or sufficient to produce anorexia in the setting of chronic inflammation.

Relatively little is known regarding the intracellular signaling mechanisms by which inflammatory mediators activate POMC neurons. Jang *et al.* [28^{••}] investigated the role of NF- κ B activation in POMC neurons during acute inflammatory challenges. LPS, IL-1 β , and TNF- α stimulate transcription of the POMC gene *in vitro*; inhibiting NF- κ B expression or activity attenuates or abolishes this effect. Transgenic mice that lack I κ B kinase- β (an enzyme upstream of NF- κ B activation) specifically in POMC neurons exhibit blunted anorexia and less weight loss in response to intra-hypothalamic LPS injection. Based on these observations, we can conclude that acute inflammation-induced anorexia and weight loss depend upon NF- κ B-mediated transcription in hypothalamic POMC neurons. Although the authors of this study focused on acute illness in their experiments, it would be interesting to ascertain whether mice with impaired NF- κ B signaling in POMC neurons resist developing

anorexia/cachexia when exposed to a chronic inflammatory insult.

Genetic or pharmacological blockade of MC4-R signaling ameliorates cachexia in many animal disease models, including acute LPS administration, cancer, and renal failure [29–31]. Scarlett *et al.* [32[•]] extended these findings to two experimental models of cardiac cachexia. Unlike sham-operated controls, wild-type mice that underwent surgery to induce myocardial infarction (MI) failed to gain body weight, lean body mass or fat mass. In contrast, mice with a targeted deletion of the MC4-R gene gained body weight, lean body mass and fat mass regardless of whether they were in the sham-operated or MI groups. Furthermore, MI reduced food intake and increased oxygen consumption in wild-type mice but not in MC4-R null mice. In an aortic banded rat model of pressure overload cardiac hypertrophy, ICV AgRP treatment for 2 weeks increased body weight, lean body mass, fat mass, and food intake compared to baseline values. The observation that chronic blockade of central MC4-R signaling protects animals from developing cardiac cachexia suggests that MC4-R antagonism might be a viable therapeutic strategy for patients with chronic heart failure (CHF). Because preexisting cachexia predicts mortality following heart transplantation, pharmacological MC4-R antagonism might be an effective pretransplant therapy [33].

Preclinical studies utilizing small molecule, peripherally administered MC4-R antagonists have yielded favorable results with respect to preventing or reversing anorexia and loss of lean mass in experimental models of acute and chronic illness [34]. Recently, MC4-R antagonists with oral bioavailability have been developed and tested in cachectic animals [35]. Weyermann *et al.* [36^{••}] reported that SNT207707 and SNT207858 are novel, nonpeptide, selective MC4-R antagonists that penetrate the blood-brain barrier. Oral administration of these compounds dose-dependently increases food intake in healthy mice. In mice implanted with a C26 adenocarcinoma, once daily oral administration of SNT207707 and SNT207858 for 2 weeks prevents tumor-induced loss of body weight, fat mass, and lean body mass without affecting tumor size. It is unknown whether the preservation of body weight in the tumor-bearing mice was solely attributable to the orexigenic activity of the MC4-R antagonists, or whether SNT207707 and SNT207858 also alter metabolism and/or energy partitioning.

Monoclonal antibody therapy is another potential anti-cachexia therapeutic strategy. mAb 1E8a is selective for the human MC4-R and behaves as an inverse agonist and noncompetitive antagonist *in vitro* [37[•]]. Acute ICV injection of purified mAb 1E8a increases body weight and food intake in rats. Chronic (7 day) ICV infusion of

mAb 1E8a stimulates food consumption, reduces weight loss, and increases adiposity. A single chain variant of mAb 1E8a that crosses the blood–brain barrier is effective at increasing food intake when administered intravenously.

To date, there have not been any published clinical trials with drugs that antagonize or impair MC4-R signaling. Given the recent promising results obtained from preclinical studies, it is presumably only a matter of time before the safety and efficacy of MC4-R antagonists are tested in humans.

Neuropeptide Y

NPY is an orexigenic peptide that is co-expressed in the same neurons as AgRP in the ARC [38]. Like AgRP, NPY mRNA expression increases during states of energy deficiency (e.g., starvation), and centrally administered NPY promotes feeding and weight gain [39]. NPY gene expression is usually either unchanged or increased in experimental models of chronic inflammation. For example, in a rat model of adjuvant arthritis, NPY mRNA levels were elevated despite reduced food intake, body weight, and adiposity compared with control animals [40]. Repeated intrahypothalamic NPY injections do not increase food intake in tumor-bearing rats, indicating desensitization to NPY in these animals [41]. Although some degree of dysfunction in the NPY system probably occurs in cachectic humans, cachexia does not appear to be an NPY-deficient state, and thus NPY replacement is unlikely to prevent or reverse cachexia in a clinical setting. Consequently, interest in NPY as an anti-cachexia therapy has waned over the past several years.

Ghrelin

Ghrelin is an orexigenic/anabolic peptide hormone that was identified by virtue of its ability to stimulate growth hormone (GH) secretion [42]. Ghrelin, which is primarily produced by neuroendocrine cells of the gastric fundus, is an agonist at the growth hormone secretagogue-1a (GHS-1a) receptor. Circulating ghrelin exists in two forms: acylated ghrelin (the biologically active form that binds and activates GHS-1a receptors) and des-acyl ghrelin (an inactive form that lacks biological activity at GHS-1a receptors) [43]. Ghrelin is a meal initiation signal, with its plasma levels rising in the fasted state and decreasing postprandially. Acute intravenous ghrelin administration increases caloric intake in humans [44], and chronic ghrelin treatment increases adiposity in mice [45]. NPY/AgRP neurons in the ARC express GHS-1a receptors [46], and both neuropeptides play essential roles in ghrelin-induced food intake [47]. In addition to its effects on energy homeostasis, ghrelin also exerts potent anti-inflammatory actions, at least in part, by decreasing

peripheral immune cell expression of pro-inflammatory cytokines [48].

Although one recent study reported reduced plasma ghrelin concentrations in cancer patients [49], cachexia is more frequently associated with increased circulating ghrelin levels compared with healthy controls or non-cachectic patients with the same underlying diseases [50–53]. In rats, plasma ghrelin concentrations are transiently reduced following acute LPS administration but are increased by repeated LPS injections [54]. In the context of cachexia, increased plasma ghrelin concentrations could represent a compensatory orexigenic/anabolic response to tissue wasting. Alternatively, elevated ghrelin concentrations in cachectic individuals might signify a state of ghrelin resistance. Many human experiments only report total circulating ghrelin levels (i.e., the investigators do not distinguish between acylated vs. des-acyl ghrelin), and thus it is difficult to draw meaningful conclusions about levels of bioactive ghrelin in those patients. Nevertheless, the bulk of the evidence suggests that cachexia is not a ghrelin-deficient state.

In rodent models of cancer, kidney disease, and CHF, administration of ghrelin or GHS-1a receptor agonists ameliorates many of the symptoms of cachexia [55–57]. In a rat model of CHF, repeated injections or chronic infusion of human ghrelin for 4 weeks elicited dose-dependent stimulatory effects on feeding, body weight gain, adiposity, and lean body mass accumulation, in the absence of any significant improvement in cardiovascular function [58[•]]. Skeletal muscle mitochondrial oxidative capacity is impaired in rats with chronic kidney disease but is corrected by ghrelin treatment, even in the absence of ghrelin-induced increases in food intake [59[•]].

Early clinical trials with ghrelin or synthetic ghrelin mimetics have shown increased caloric intake, body weight, and lean body mass in cachectic patients with various cancers, chronic kidney disease, congestive heart failure, and chronic obstructive pulmonary disorder [60]. Ghrelin has a short half-life (30 min) in the circulation and must be injected, both of which limit its utility as a therapeutic agent. Phase I and II clinical trial with the orally available, synthetic ghrelin mimetic RC-1291 have demonstrated increases in body weight and lean body mass in healthy individuals [61] and in cancer patients [62], without any dose-limiting side effects. Ghrelin's ability to stimulate the secretion of GH (and thus, insulin-like growth factor-1) raises the concern that ghrelin mimetics might increase tumor growth in cancer patients. To date, there have not been any published randomized, placebo-controlled clinical trials investigating outcomes of long-term treatment with ghrelin

analogues in cachectic patients. Until these trials are performed, we can only speculate as to whether long-term administration of ghrelin analogues would elicit sustained increases in caloric intake and body mass accrual in cachectic individuals.

Orexins/hypocretins

Orexin (also known as hypocretin) neurons in the lateral hypothalamus regulate feeding behavior, arousal, and motivated behaviors. Central administration of orexin-A acutely stimulates feeding but does not elicit sustained increases in feeding or body weight during chronic administration in rats [63,64]. Injection of orexin-A into the ventromedial hypothalamus stimulates glucose metabolism and increases insulin sensitivity in skeletal muscle via activation of the sympathetic nervous system [65^{••}]. Because orexins promote wakefulness and increase locomotor activity [66], it is conceivable that a sustained reduction in orexin neuronal activity might underlie lethargy in cachectic individuals. However, the response of orexin neurons to an acute inflammatory challenge is equivocal. In rats, peripherally administered LPS either increases or decreases Fos expression in orexin neurons during the dark phase of the light/dark cycle, which is when orexin neurons are maximally activated [67[•],68]. In pilot studies, the psychostimulant modafinil (which stimulates orexin neurons) reduced fatigue in cancer patients [69[•],70[•]]. A potential role for orexins in the pathophysiology of cachexia remains largely unexplored.

Conclusion

Cachexia is characterized by a profound dysregulation of metabolic function. The neural mechanisms contributing to the pathophysiology of cachexia are complex, and as such remain poorly defined. Well tolerated and effective pharmacological treatments for cachexia are currently lacking, but randomized, placebo-controlled clinical trials with MC4-R antagonists or long-term treatment with ghrelin mimetics are eagerly anticipated. Although most researchers in the cachexia field study muscle catabolism and/or appetite regulation, it is important not to ignore the negative impact of other cachexia-related symptoms (e.g., lethargy). Elucidating the mechanisms by which inflammatory mediators promote muscle wasting, anorexia, and other illness behaviors will not only shed light upon the basic biology of cachexia and its underlying disease processes, but also hopefully identify novel therapeutic strategies for treating cachexia of various etiologies.

Acknowledgements

The authors do not have any conflicts of interest to declare. This work was supported by grants from the National Institutes of Health (NIDDK).

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

- 1 Tisdale MJ. Cancer cachexia. *Curr Opin Gastroenterol* 2010; 26:146–151.
- 2 Cheung WW, Paik KH, Mak RH. Inflammation and cachexia in chronic kidney disease. *Pediatr Nephrol* 2010; 25:711–724.
- 3 Freeman LM. The pathophysiology of cardiac cachexia. *Curr Opin Support Palliat Care* 2009; 3:276–281.
- 4 Faintuch J, Soeters PB, Osmo HG. Nutritional and metabolic abnormalities in pre-AIDS HIV infection. *Nutrition* 2006; 22:683–690.
- 5 Evans WJ, Morley JE, Argiles J, *et al.* Cachexia: a new definition. *Clin Nutr* 2008; 27:793–799.
- 6 Evans WK, Makuch R, Clamon GH, *et al.* Limited impact of total parenteral nutrition on nutritional status during treatment for small cell lung cancer. *Cancer Res* 1985; 45:3347–3353.
- 7 Morley JE, Thomas DR, Wilson MM. Cachexia: pathophysiology and clinical relevance. *Am J Clin Nutr* 2006; 83:735–743.
- 8 Mak RH, Cheung W, Cone RD, Marks DL. Mechanisms of disease: cytokine and adipokine signaling in uremic cachexia. *Nat Clin Pract Nephrol* 2006; 2:527–534.
- 9 Ren SG, Seliktar J, Li X, *et al.* Measurement of leukemia inhibitory factor in biological fluids by radioimmunoassay. *J Clin Endocrinol Metab* 1998; 83:1275–1283.
- 10 Grunfeld C, Feingold KR. Regulation of lipid metabolism by cytokines during host defense. *Nutrition* 1996; 12:S24–S26.
- 11 Acharyya S, Guttridge DC. Cancer cachexia signaling pathways continue to emerge yet much still points to the proteasome. *Clin Cancer Res* 2007; 13:1356–1361.
- 12 Layé S, Parnet P, Goujon E, Dantzer R. Peripheral administration of lipopolysaccharide induces the expression of cytokine transcripts in the brain and pituitary of mice. *Brain Res Mol Brain Res* 1994; 27:157–162.
- 13 Grossberg AJ, Scarlett JM, Marks DL. Hypothalamic mechanisms in cachexia. • *Physiol Behav* 2010; 100:478–489.
A comprehensive review on hypothalamic neuropeptides that mediate illness-induced anorexia and lesser studied aspects of cachexia, including lethargy, insulin resistance and reproductive dysfunction.
- 14 Laflamme N, Rivest S. Effects of systemic immunogenic insults and circulating proinflammatory cytokines on the transcription of the inhibitory factor kappaB alpha within specific cellular populations of the rat brain. *J Neurochem* 1999; 73:309–321.
- 15 Elmquist JK, Scammell TE, Jacobson CD, Saper CB. Distribution of Fos-like immunoreactivity in the rat brain following intravenous lipopolysaccharide administration. *J Comp Neurol* 1996; 371:85–103.
- 16 Wisse BE, Ogimoto K, Tang J, *et al.* Evidence that lipopolysaccharide-induced anorexia depends upon central, rather than peripheral, inflammatory signals. *Endocrinology* 2007; 148:5230–5237.
- 17 Plata-Salamán CR. Anorexia induced by activators of the signal transducer gp 130. *Neuroreport* 1996; 7:841–844.
- 18 Sonti G, Ilyin SE, Plata-Salamán CR. Anorexia induced by cytokine interactions at pathophysiological concentrations. *Am J Physiol* 1996; 270:R1394–R1402.
- 19 Wallenius K, Wallenius V, Sunter D, *et al.* Intracerebroventricular interleukin-6 treatment decreases body fat in rats. *Biochem Biophys Res Commun* 2002; 293:560–565.
- 20 DeBoer MD, Scarlett JM, Levasseur PR, *et al.* Administration of IL-1 β to the 4th ventricle causes anorexia that is blocked by agouti-related peptide and that coincides with activation of tyrosine-hydroxylase neurons in the nucleus of the solitary tract. *Peptides* 2009; 30:210–218.
- 21 Arruda AP, Milanski M, Romanatto T, *et al.* Hypothalamic actions of tumor necrosis factor alpha provide the thermogenic core for the wastage syndrome in cachexia. *Endocrinology* 2010; 151:683–694.
This paper demonstrates an essential role for central TNF- α signaling in two rodent models of cachexia.
- 22 Tao YX. The melanocortin-4 receptor: physiology, pharmacology, and pathophysiology. *Endocr Rev* 2010. [Epub ahead of print]
- 23 Ollmann MM, Wilson BD, Yang YK, *et al.* Antagonism of central melanocortin receptors in vitro and in vivo by agouti-related protein. *Science* 1997; 278:135–138.
- 24 Scarlett JM, Jobst EE, Enriori PJ, *et al.* Regulation of central melanocortin signaling by interleukin-1 beta. *Endocrinology* 2007; 148:4217–4225.
- 25 Scarlett JM, Zhu X, Enriori PJ, *et al.* Regulation of agouti-related protein messenger ribonucleic acid transcription and peptide secretion by acute and chronic inflammation. *Endocrinology* 2008; 149:4837–4845.
- 26 Macia L, Viltart O, Delacore M, *et al.* Interleukin-7, a new cytokine targeting the mouse hypothalamic arcuate nucleus: role in body weight and food intake regulation. *PLoS One* 2010; 5:e9953.
This paper reveals a novel role for interleukin-7 in the central control of feeding and body weight regulation, possibly via its actions on hypothalamic POMC and AgRP neurons.
- 27 Grossberg AJ, Scarlett JM, Zhu X, *et al.* Arcuate nucleus proopiomelanocortin •• neurons mediate the acute anorectic actions of leukemia inhibitory factor via gp130. *Endocrinology* 2010; 151:606–616.
This series of experiments demonstrated that POMC neurons are direct targets for central LIF signaling, and that gp130 signaling in POMC neurons is essential for acute LIF-induced anorexia in mice.
- 28 Jang PG, Namkoong C, Kang GM, *et al.* NF-kappaB activation in hypothalamic •• pro-opiomelanocortin neurons is essential in illness- and leptin-induced anorexia. *J Biol Chem* 2010; 285:9706–9715.
This study is among the first to investigate the intracellular signaling mechanisms by which inflammatory mediators activate POMC neurons in the setting of acute illness/inflammation. Notably, mice with impaired NF- κ B function in POMC neurons resist developing anorexia in response to acute inflammatory challenges.
- 29 Cheung W, Yu PX, Little BM, *et al.* Role of leptin and melanocortin signaling in uremia-associated cachexia. *J Clin Invest* 2005; 115:1659–1665.
- 30 Marks DL, Ling N, Cone RD. Role of the central melanocortin system in cachexia. *Cancer Res* 2001; 61:1432–1438.
- 31 Wisse BE, Frayo RS, Schwartz MW, Cummings DE. Reversal of cancer anorexia by blockade of central melanocortin receptors in rats. *Endocrinology* 2001; 142:3292–3301.
- 32 Scarlett JM, Bowe DD, Zhu X, *et al.* Genetic and pharmacologic blockade of • central melanocortin signaling attenuates cardiac cachexia in rodent models of heart failure. *J Endocrinol* 2010; 206:121–130.
These experiments revealed the beneficial effects of melanocortin receptor blockade on feeding, weight gain and lean body mass accumulation in two experimental models of cardiac cachexia.
- 33 Lietz K, John R, Burke EA, *et al.* Pretransplant cachexia and morbid obesity are predictors of increased mortality after heart transplantation. *Transplantation* 2001; 72:277–283.
- 34 DeBoer MD. Update on melanocortin interventions for cachexia: progress toward clinical application. *Nutrition* 2010; 26:146–151.
- 35 Chen C, Tucci FC, Jiang W, *et al.* Pharmacological and pharmacokinetic characterization of 2-piperazine-alpha-isopropyl benzylamine derivatives as melanocortin-4 receptor antagonists. *Bioorg Med Chem* 2008; 16:5606–5618.
- 36 Weyermann P, Dallmann R, Magyar J, *et al.* Orally available selective melanocortin-4 receptor antagonists stimulate food intake and reduce cancer-induced cachexia in mice. *PLoS One* 2009; 4:e4774.
One of the first published preclinical studies investigating the efficacy of orally available melanocortin receptor antagonists. Both compounds attenuated the loss of lean and fat mass in tumor-bearing mice.
- 37 Peter JC, Lecourt AC, Weckering M, *et al.* A pharmacologically active • monoclonal antibody against the human melanocortin-4 receptor: effectiveness after peripheral and central administration. *J Pharmacol Exp Ther* 2010; 333:478–490.
Results from these experiments in healthy rats support the idea that monoclonal antibody therapy may be a viable anti-cachexia therapy.
- 38 Hahn TM, Breininger JF, Baskin DG, Schwartz MW. Coexpression of AgRP and NPY in fasting-activated hypothalamic neurons. *Nat Neurosci* 1998; 1:271–272.
- 39 Chee MJ, Colmers WF. Y eat? *Nutrition* 2008; 24:869–877.
- 40 Stofkova A, Haluzik M, Zelezna B, *et al.* Enhanced expressions of mRNA for neuropeptide Y and interleukin 1 beta in hypothalamic arcuate nuclei during adjuvant arthritis-induced anorexia in Lewis rats. *Neuroimmunomodulation* 2009; 16:377–384.
- 41 Chance WT, Balasubramanian A, Thompson H, *et al.* Assessment of feeding response of tumor-bearing rats to hypothalamic injection and infusion of neuropeptide Y. *Peptides* 1996; 17:797–801.
- 42 Kojima M, Hosoda H, Date Y, *et al.* Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* 1999; 402:656–660.
- 43 Castaneda TR, Tong J, Datta R, *et al.* Ghrelin in the regulation of body weight and metabolism. *Front Neuroendocrinol* 2010; 31:44–60.

- 44 Akamizu T, Takaya K, Irako T, *et al.* Pharmacokinetics, safety, and endocrine and appetite effects of ghrelin administration in young healthy subjects. *Eur J Endocrinol* 2004; 150:447–455.
- 45 Tschöp M, Smiley DL, Heiman ML. Ghrelin induces adiposity in rodents. *Nature* 2000; 407:908–913.
- 46 Willesen MG, Kristensen P, Rømer J. Co-localization of growth hormone secretagogue receptor and NPY mRNA in the arcuate nucleus of the rat. *Neuroendocrinology* 1999; 70:306–316.
- 47 Chen HY, Trumbauer ME, Chen AS, *et al.* Orexigenic action of peripheral ghrelin is mediated by neuropeptide Y and agouti-related protein. *Endocrinology* 2004; 145:2607–2612.
- 48 Dixit VD, Schaffer EM, Pyle RS, *et al.* Ghrelin inhibits leptin- and activation-induced proinflammatory cytokine expression by human monocytes and T cells. *J Clin Invest* 2004; 114:57–66.
- 49 Legakis I, Stathopoulos J, Matzouridis T, Stathopoulos GP. Decreased plasma ghrelin levels in patients with advanced cancer and weight loss in comparison to healthy individuals. *Anticancer Res* 2009; 29:3949–3952.
- 50 Takahashi M, Terashima M, Takagane A, *et al.* Ghrelin and leptin levels in cachectic patients with cancer of the digestive organs. *Int J Clin Oncol* 2009; 14:315–320.
- 51 Nagaya N, Uematsu M, Kojima M, *et al.* Elevated circulating level of ghrelin in cachexia associated with chronic heart failure: relationships between ghrelin and anabolic/catabolic factors. *Circulation* 2001; 104:2034–2038.
- 52 Yoshimoto A, Mori K, Sugawara A, *et al.* Plasma ghrelin and desacyl ghrelin concentrations in renal failure. *J Am Soc Nephrol* 2002; 13:2748–2752.
- 53 Itoh T, Nagaya N, Yoshikawa M, *et al.* Elevated plasma ghrelin level in underweight patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004; 170:879–882.
- 54 Hataya Y, Akamizu T, Hosoda H, *et al.* Alterations of plasma ghrelin levels in rats with lipopolysaccharide-induced wasting syndrome and effects of ghrelin treatment on the syndrome. *Endocrinology* 2003; 144:5365–5371.
- 55 DeBoer MD, Zhu X, Levasseur PR, *et al.* Ghrelin treatment of chronic kidney disease: improvements in lean body mass and cytokine profile. *Endocrinology* 2008; 149:827–835.
- 56 DeBoer MD, Zhu XX, Levasseur P, *et al.* Ghrelin treatment causes increased food intake and retention of lean body mass in a rat model of cancer cachexia. *Endocrinology* 2007; 148:3004–3012.
- 57 Nagaya N, Uematsu M, Kojima M, *et al.* Chronic administration of ghrelin improves left ventricular dysfunction and attenuates development of cardiac cachexia in rats with heart failure. *Circulation* 2001; 104:1430–1435.
- 58 Akashi YJ, Palus S, Datta R, *et al.* No effects of human ghrelin on cardiac function despite profound effects on body composition in a rat model of heart failure. *Int J Cardiol* 2009; 137:267–275.
- These experiments demonstrated that chronic ghrelin treatment improves lean body mass retention in a rat model of chronic heart failure, in the absence of any significant improvements in cardiac function.
- 59 Barazzoni R, Zhu X, Deboer M, *et al.* Combined effects of ghrelin and higher food intake enhance skeletal muscle mitochondrial oxidative capacity and AKT phosphorylation in rats with chronic kidney disease. *Kidney Int* 2010; 77:23–28.
- Exogenous ghrelin treatment improves metabolic function in the skeletal muscle of nephrectomized rats, at least in part via mechanisms that operate independently of ghrelin-induced food intake.
- 60 Gullett NP, Hebbar G, Ziegler TR. Update on clinical trials of growth factors and anabolic steroids in cachexia and wasting. *Am J Clin Nutr* 2010; 91:1143S–1147S.
- 61 Garcia JM, Polvino WJ. Effect on body weight and safety of RC-1291, a novel, orally available ghrelin mimetic and growth hormone secretagogue: results of a phase I, randomized, placebo-controlled, multiple-dose study in healthy volunteers. *Oncologist* 2007; 12:594–600.
- 62 Garcia JM, Graham C, Kumor K, Polvino W. A Phase II, randomized, placebo-controlled, double blind study of the efficacy and safety of RC-1291 for the treatment of cancer-cachexia. *J Clin Oncol* 2007; 25:9133 [abstract].
- 63 Haynes AC, Jackson B, Overend P, *et al.* Effects of single and chronic intracerebroventricular administration of the orexins on feeding in the rat. *Peptides* 1999; 20:1099–1105.
- 64 Yamanaka A, Sakurai T, Katsumoto T, *et al.* Chronic intracerebroventricular administration of orexin-A to rats increases food intake in daytime, but has no effect on body weight. *Brain Res* 1999; 849:248–252.
- 65 Shiuchi T, Haque MS, Okamoto S, *et al.* Hypothalamic orexin stimulates feeding-associated glucose utilization in skeletal muscle via sympathetic nervous system. *Cell Metab* 2009; 10:466–480.
- This paper reveals a role for hypothalamic orexin neurons in skeletal muscle glucose metabolism.
- 66 Anaclet C, Parmentier R, Ouk K, *et al.* Orexin/hypocretin and histamine: distinct roles in the control of wakefulness demonstrated using knock-out mouse models. *J Neurosci* 2009; 29:14423–14438.
- 67 Gaykema RP, Goehler LE. Lipopolysaccharide challenge-induced suppression of Fos in hypothalamic orexin neurons: their potential role in sickness behavior. *Brain Behav Immun* 2009; 23:926–930.
- These experiments hint at a potential role of reduced activity of hypothalamic orexin neurons in illness-induced lethargy.
- 68 Hollis JH, Lemus M, Evetts MJ, Oldfield BJ. Central interleukin-10 attenuates lipopolysaccharide-induced changes in food intake, energy expenditure and hypothalamic Fos expression. *Neuropharmacology* 2010; 58:730–738.
- 69 Blackhall L, Petroni G, Shu J, *et al.* A pilot study evaluating the safety and efficacy of modafinil for cancer-related fatigue. *J Palliat Med* 2009; 12:433–439.
- This small study in human subjects demonstrated that modafinil (a CNS stimulant that increases orexin activity) reduces fatigue and improves quality of life in cancer patients.
- 70 Spathis A, Dhillan R, Booden D, *et al.* Modafinil for the treatment of fatigue in lung cancer: a pilot study. *Palliat Med* 2009; 23:325–331.
- Another pilot study that tested the safety and efficacy of modafinil in cancer patients. Modafinil lessened fatigue, perhaps by virtue of its ability to stimulate hypothalamic orexin neuronal activity.